



Significant reductions in oil quality and lipid content of oilseed rape (*Brassica napus* L.) under climate change

Namazkar, Shahla; Egsgaard, Helge; Frenck, Georg ; Terkelsen, Thilde ; Bagger Jørgensen, Rikke

Published in:
Procedia Environmental Sciences

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Namazkar, S., Egsgaard, H., Frenck, G., Terkelsen, T., & Bagger Jørgensen, R. (2015). Significant reductions in oil quality and lipid content of oilseed rape (*Brassica napus* L.) under climate change. *Procedia Environmental Sciences*, 29, 121-122.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Agriculture and Climate Change - Adapting Crops to Increased Uncertainty (AGRI 2015)

Significant reductions in oil quality and lipid content of oilseed rape (*Brassica napus* L.) under climate change

Shahla Namazkar^a, Helge Egsgaard^a, Georg Frenck^b, Thilde Terkelsen^c and Rikke Bagger Jørgensen^{a*}

^a Department of Chemical and Biochemical Engineering, Technical University of Denmark, Risø Campus, Frederiksborgvej 399, DK-4000, Roskilde, Denmark

^b University of Innsbruck, Institute of Ecology, Sternwartestraße 15, 6020 Innsbruck, Austria

^c University of Copenhagen, Copenhagen Biocenter, Ole Maaløes Vej 5, 2200 København N, Denmark

*Corresponding author rijq@kt.dtu.dk

Abstract

Despite of the potential importance to food and bioenergy purposes, effects from climate change on plant oil quality have hardly been characterized.

On a global basis *Brassica napus* L., rapeseed or oilseed rape, is the second largest source of vegetable oil after soybean and the predominant oil crop in Europe. We found significant changes in oil quality and quantity of four cultivars of oilseed rape grown in five future climate scenarios with elevated [CO₂], [O₃], temperature and combinations hereof (~RCP8.5₁(1)). Populations of the cultivars were grown under ambient and climate change conditions in a climate-phytotron. The treatments were ambient (360 ppm CO₂, 19/12 °C (day/night), 20/20 ppb O₃ (day/night)), all factors elevated (650 ppm CO₂, 24/17 °C, 60/20 ppb O₃), as well as two- and single-factor treatments with the elevated factors.

The overall trend was that oil content and quality were significantly reduced, except in the scenario with elevated [CO₂] alone. Of the six analyzed fatty acids five - oleic acid (C18:1), linoleic acid (C18:2), linolenic acid (C18:3, omega-3), palmitic (C16:0), eicosenoic acid (C20:1) - showed reductions, the only exception being stearic acid, C18:0. For example we found that in the two-factor treatment, where elevated [CO₂] and temperature were combined, the essential fatty acid omega-3, C18:3, decreased by 45% and oil content declined 10%.

Total losses in fatty acid and oil yields would be even larger, when also considering reported reductions in seed biomass in the future scenarios (2,3): We estimate that when [CO₂] and temperature are elevated simultaneously, the oil yield per hectare will drop 58% and the production of omega-3 (C18:3) will be reduced by 77%/hectare. Also the proportion between saturated and unsaturated fatty acids was changed for the worse. Facing this outlook, breeding for climate tolerant cultivars seems essential for oil yield and quality.

© 2015 Published by Elsevier B.V This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Peer-review under responsibility of the organizing committee of the Agriculture and Climate Change - Adapting Crops to Increased Uncertainty (AGRI 2015)

Keywords: Climate change; oilseed rape quality; oil quantity; fatty acids; oleic acid linoleic acid; linolenic acid; palmitic acid; eicosenoic acid; stearic acid

References

1. Stocker TF *et al.* IPCC. Summary for Policymakers. In: Climate Change 2013: *The Physical Science Basis*. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. 2013
2. Frencik G, Linden LD, Mikkelsen TN, Brix H, Jørgensen RB. Increased [CO₂] does not compensate for negative effects on yield caused by higher temperature and [O₃] in *Brassica napus* L. *Eur. J. Agron.* 2011; **35**, 127– 134.
3. Frencik G, Linden L, Mikkelsen TN, Brix H, Jørgensen RB. Response to multi-generational selection under elevated [CO₂] in two temperature regimes suggests enhanced carbon assimilation and increased reproductive output in *Brassica napus* L. *Ecol. Evol.* 2013; **3**, 1163–1172 (2013).